

## The Effects of Sounds and Music on Cells and Organisms: A Promising and Developing Area of Research

By Claire Brun\* & Jean-Marie Exbrayat<sup>‡</sup>

*This review is based on recent publications about the effects of sound, and more particularly of music, on several aspects of physiology. It has been known for a long time that music has effects on the brain and on the functioning of different organs. In recent years, several publications also described specific effects of music on the physicochemical mechanisms in the other organisms, bacteria, plants and animals. These researches being rather disparate in the methodologies used and the results obtained, they need to be classified. In this review, we summarize the studies and attempt to explain the cellular mechanisms involved, by considering the properties of the plasma membrane and its links with the extracellular and intracellular medium. This field of research is currently in full expansion, but still requires further studies to understand and go further in the possible applications, the precise molecular mechanisms of effects of music still remain to be clarified.*

**Keywords:** *sound, music, plant, animal, unicellular organism*

### Introduction

Music is a complex acoustic and temporal structure, whose effects on biology are a hot topic. It has long been known that music influences mood and arouses strong emotions. Today, we are discovering the numerous effects it has on the brain and on organic functions. It is used in a care approach (music therapy) (Behzadmehr et al. 2020, Billar et al. 2020, Buglione et al. 2020, Bulut et al. 2020, Martin et al. 2020, Çelebi et al. 2020, Chai et al. 2020a, b, Cimen et al. 2020, Chan et al. 2020, Dai et al. 2020, Ernberg et al. 2020, Gamboa et al. 2020, Garcia Guerra et al. 2020, Giordano et al. 2020, Gogoularadja and Bakshi 2020, Guo et al. 2020, Howlin and Rooney 2020, Johnson and Elkins 2020, Köhler et al. 2020, Pérez-Eizaguirre and Vergara-Moragues 2020, Polascik et al. 2021, Saraogi et al. 2020, Usui et al. 2020, Wang et al. 2020, Zhang et al. 2020, Dunbar et al. 2021, Yang et al. 2021), but its influence on well-being goes beyond the human being (Sambraus and Hecker 1985, Hurnik and Johnson 1997, Kenison 2016, Bowman et al. 2017, González-Grajales et al. 2019, Kemp 2020). In fact, many studies have described the various effects of music on different organisms (Lemarquis 2009, 2021, Dhungana et al. 2018, Exbrayat and Brun 2019, Mayoud and Lemarquis 2019, Pulcrano and di Porzio 2021).

---

\*Professor, Lyon Catholic University, France.

<sup>‡</sup>Emeritus Professor, Lyon Catholic University, and Ecole Pratique des Hautes Etudes, France.

Music is a fundamental mean of communication, which developed and diversified over the millennia as human beings migrated around the planet (Wallin et al. 2000, Mehr et al. 2020, Savage et al. 2020). It allows people to share intentions and emotions. Power of music pass through the capture of musical sounds by auditory cells which record sounds, transform them into electrical signals which in turn reach the brain where they are translated into emotions. The neurophysiological mechanisms are better known thanks to the development of innovated methods (Zimmermann and Fermin 1996, Fettiplace and Hackney 2006, Ashmore 2008, Müller 2008, Rabbitt et al. 2010). Scientific evidences thus confirmed the long-praised medicinal virtues of music, such as neurostimulation and neuroprotection (Lemarquis 2009, 2021, Jones et al. 2020). But for a few years now, more and more publications have addressed the direct effects on non-auditory cells, a less obvious aspect of the effects of music (Lestard et al. 2013, Lestard and Capella 2016). The experiences performed by researchers present very disparate results. Whatever this great disparity, the study of the effects of sound and music on cells reveals the available possibilities to modulate cell physiology.

### **Evolution and Adaptation of Communication in the Animal Kingdom**

So that an organism could live, it is necessary that it communicates with other organisms and, in a broader sense, with the elements of the environment. Throughout its life, every organism is subject to many stimuli. In unicellular organisms, the single cell is equipped with receptors that register different types of stimuli acting directly on it. Some unicellular, prokaryotes or eukaryotes, are sensitive to molecules emitted by other individuals and react by synthesizing substances. They may also be sensitive to vibrations produced by sound and respond by producing molecules. For example, among unicellular organisms, volvox algae live in colonies of thousands of flagellated cells, able to move toward the same direction. All the cells contain a light-sensitive pigment that directs each cell towards it, and the perception of light allows a set of juxtaposed cells to move towards the same goal (Hallmann 2003, Ikushima and Maruyama 2007).

In sponges, jellyfish and polyps, a reduced number of cell types is observed and the nervous system is limited to a few neurons, or it is almost absent (Arai 1997, Moore 2006, Barnes et al. 2009). The perception of external signals exists however and allows oriented movements and responses adapted to the environment. As organisms become more complex, various sensory organs appear, in connection with the development of an increasingly efficient nervous system. The latter will allow a better adaptation of the organisms in their environment. Stimuli are then received by receptors of sense organs, and the nerve impulses are transmitted and interpreted by the nervous system.

For Humans, the five specific senses commonly known and already listed by Aristotle (384-322 BC), are touch, sight, hearing, smell and taste. In addition, there is a general sensitivity including mechanoreception, thermoception, nociception and proprioception.

There are also other types of senses in the animal kingdom directly related to the evolutionary level and the way of life of organisms. For example, fish and certain amphibian larvae have a lateral line which allows perception of water movements and hydrostatic pressure (Bleckmann and Zelick 2009). They also have an ampullary organ (consisting of the ampullae described by Lorenzini in 1678) which is sensitive to low frequency electric fields and allows electrolocation (Roth and Tscharrntke 1976, Gibbs and Northcutt 2004, King et al. 2018). In lizards, snakes and caecilian amphibians, particularly well-developed vomeronasal organs are added to the classic olfactory organ (Badenhorst 1978, Billo 1986, Billo and Wake 1987, Døving and Trotier 1998). Finally, in some migratory birds, magnetoreceptors make the individual sensitive to the earth's magnetic field (Wiltschko and Wiltschko 2012).

### **Sounds and Music as a Communication Signal**

In the course of evolution, different mechanisms have been selected to allow animals to emit sounds and communicate.

Aristotle already indicated that fish could emit “vocalizations”. A significant number of studies confirm that some fish emit sounds and that their repertoire is surprisingly vast. So, sound is involved in a large number of behaviors related to reproduction, feeding or defense of the territory. The means of producing sounds are variable; they range from the air bubble released by the posterior part of digestive tube (in herrings) to chirping produced by the friction of two hard parts of the body: striated joint of the pectoral fins in catfish, pharyngeal teeth located at the level of the gills which squeak in *Haemulon plumierii*; some species can also use the swim bladder to emit sounds (Parmentier et al. 2016, Raick et al. 2018, Di Iorio et al. 2019, Huby et al. 2019, Bolgan et al. 2020). About 100 families of bony fish have been shown to be able to communicate acoustically. The same species can even produce several different sounds. Currently, the "vocalizations" of more than 200 species of fish were described in the Northwest Atlantic (Fish and Mowbray 1970).

Aquatic mammals such as Cetaceans also emit all kinds of sounds, vocally or not, allowing communication between animals of the same species (Payne 1984). Sonar is used by dolphins (Whitlow 1993). These animals emit sounds of varying frequencies to communicate or find their way in space. They use air sacs (Helmholtz cavities) to emit sounds at various resonant frequencies. When diving, dolphins store air in the lungs. They use their larynx to produce ultrasounds (Whitlow 1993).

Bats use echolocation to navigate and locate prey. They modulate calls of different frequencies according to the species, which allow them to perceive the distance of the prey or the obstacle by measuring the delay between emission of the call and echo. The perception of sound is carried out by a classical hearing system (Suga and O'Neill 1979, Moss and Sinha 2003).

In many species of birds, at the time of reproduction, the male utters a characteristic song using the syrinx, an organ made up of cartilaginous parts

located below the trachea, unlike the vocal cords which are located above the trachea (Warner 2009).

These few examples show that the senses of organisms represent adaptations to living environment. Any response to the environment implies that stimuli such as molecules, vibrations, low frequencies, electric fields or light waves, are captured at a membrane receptor, before a cascade of intracellular reactions leads to a response (Lim et al. 2015, Syrovatkina et al. 2016). The complexity of the sense organs observed in multicellular organisms is closely related to the development of the nervous system: from a simple network of nerve fibers in the mesoglea (mesohyle) of porifers and cnidarians, this system becomes more complex and centralized, allowing more elaborate behavioral responses. Table 1 summarizes the diversity of sound production mechanisms in animals.

**Table 1.** *Examples Illustrating the Diversity of Sound Production Mechanisms in Animals*

Structures used in sound production	Mechanisms	Animals
Use of resonant external substrate	Hitting a substrate	<b>Beavers:</b> slapping the tail on the surface of water <b>Spiders:</b> drumming during courtship sequences
Use of body structures	Clicking, snapping, vibrating	<b>Some insects:</b> wing vibrating <b>Cicadas:</b> vibrating a drum-like membrane (modified membrane located on the abdomen) <b>Fish (Sciaenidae):</b> swimbladder snapping <b>Birds:</b> beak clicking/grinding <b>Bats:</b> tongue clicking
	Rubbing body parts together	<b>Orthopterans:</b> stridulation by dragging the body part equipped with a scraper across an adjacent body part with a file-like structure <b>Fish (catfish):</b> stridulation by using the pectoral fins
	Vibrating a membrane in an air flow	<b>Birds:</b> tympaniform membranes vibrating (presence of a syrinx) <b>Amphibians, reptiles and mammals:</b> presence of a larynx containing vocal cords

### Effects of Vibrations, Sounds and Music on Cells and Organisms

In recent years, research has focused on the effects of music on various organisms in an attempt to understand how sound vibrations act at the cell level. Experiments have been carried out on cell cultures, in cancer cells, microorganisms, and on complete organisms, by testing different biological parameters (Lemarquis 2009, 2021, Dhungana, et al. 2018, Exbrayat and Brun 2019, Mayoud and Lemarquis 2019, Pulcrano and di Porzio 2021). Tables 2 and 3 summarize the state of the art about the positive effects of music in different organisms and non-auditory cells.

**Table 2.** *State of the Art About Some Effects of Music on Non-auditory Animal Cultured Cells*

Cultured cells	Effects of music
ST2 stromal cells C2C12 myoblasts	Suppression of mechanosensitive genes by audible sound stimulation
NIH3T3 fibroblasts NB2a neuroblastoma cells	No observation of suppression of mechanosensitive genes by audible sound stimulation, showing a cell type-specific response to sound
Human embryo	Improvement of fertilization rates
Cancer cell lines	Alteration of cell cycle and decrease in viability, but differences in response depending on cell line
Buffalo granulosa cell speroids	Up-regulation of steroidogenic gene expression

Note: The key references are mentioned in the text.

**Table 3.** *State of the Art About Positive Effects of Music (Soft Rhythmic Music) in Different Uni- and Pluricellular Organisms*

Organisms		Positive effects of music	
Animal species	Domestic animals	Dogs	Increase in sleep duration
		Cats	Increase in approach behaviour
	Farm animals	Cows	Increase in milk production
		Buffaloes	Change in behaviour
	Race horses		Influence of emotional state
	Birds	Chickens	Increase in number of neurons and mean neuronal nuclear area
		Songbirds	Improvement in learning to sing
	Rodents		Improvement in spatial learning and reduction of anxiety-related behavior; increase in the body weight of pups
	Monkeys	Chimpanzees	Reduction in aggression
		Baboons	Decrease in heartrate
Elephants		Reduction of stereotyped behaviour	
Vegetal species	Herbaceous plants	Increase in germination, growth and development, disease resistance	
Unicellular organisms	Bacteria and yeast	Improvement in growth and antibiotic sensitivity	

Note: Choice of music, depending of the acoustic communication system of the species, is decisive to observe behavioral effect. In general, loud and inharmonious music tends to have a negative effect.

The key references are mentioned in the text.

### *Effects of Vibration, Sound and Music on the Animal Cells*

The effects of electromagnetic waves on living beings have been studied for much longer than those of sound waves (National Research Council (US) Committee on Assessment of the Possible Health Effects of Ground Wave Emergency Network (GWEN) 1993, Cardinale and Pope 2003). Nevertheless, a number of studies suggest that emotions triggered by music can be particularly useful in relieving stress. Recent experiments show that different non-hearing cells respond to sound, the fluids contained in the cells being sensitive to pressure variations induced by sound waves. The mechanisms of cell growth or cell death affected by acoustic vibrations appear to be similar for all cell types, whether they

are auditory or not. The basic mechanisms could thus be common and universal (Zimmermann and Fermin 1996, Chan and Hudspeth 2005, Fettiplace and Hackney 2006, Ashmore 2008, Müller 2008, Rabbitt and Boyle 2010, Lestard et al. 2013, Lestard and Capella 2016). Thus, research has been carried out on the effects of music on the biology of different cell types. Study of effects of music on human chondrocytes has shown that use of music help to identify biomarkers and provide a new approach to the treatment of osteoarthritis (Vannoni 2012, Corallo et al. 2013, 2014, Dhungana et al, 2018, Exbrayat and Brun, 2019). On the other hand, it has been observed that the size and the granularity of tumoral MCF7 cells cultivated in vitro was altered by music (Lestard et al. 2013, Lestard and Capella 2016). In the lab Sprague-Dawley rats, music also decreased the enhancing effects of stress on the development of lung metastases provoked by previously injected carcinosarcoma cells (Nuñez et al. 2002).

Music has an effect on the deformability and aggregation of red blood cells and can therefore have an influence on some pathologies. In particular, effects on the surface properties of plasmic membrane have been observed. Membrane molecules such as adenylate kinase, show sensibility to the exposure to low frequency fields (Erken et al. 2008, Albanese et al. 2009). Acoustic vibrations affect fibroblast migration and cell morphology with formation of filopodia and lamellipodia (Mohammed et al. 2016).

#### *Specific Effects on the Immune System*

Several studies have shown the effects of music on the immune system of people under stress, or suffering from Alzheimer's disease, Parkinson's disease, or after a stroke (Hasegawa et al. 2001). The effects are visible on the activity of the natural killers (NK) cells and on the hormone norepinephrine level. Other researches have shown that rhythmic percussions can increase NKs cells and the amount of the hormone dehydroepiandrosterone relative to cortisol, indicating stress regulation (Lu et al. 2013). A variety of neuroendocrine ways also contribute to immune system changes (Bittman et al 2001, Hirokawa and Ohira 2003, Wachi et al, 2007).

Several experiments have also been carried out in rodents with pathologies and have shown a role for music on the immune system. In young and adult rats suffering from asthma, music modulates the number of leukocytes and the level of IL-4 (Lu et al. 2010). In lab BALB/c mice subjected either to noise or to music, thymus and spleen cell density, T cell population, splenocyte proliferation and NKs activity were enhanced by music (Nuñez et al. 2002). The survival of allografts in an experimental model of murine heart transplantation was significantly prolonged in animals exposed to occidental opera and classical music. Cell proliferation, IL-2 and interferon- $\gamma$  were suppressed in mice exposed to opera, while IL-4 and IL-10 were upregulated and CD4 +, CD25 +, Foxp3 + increased after exposure to certain types of music (Uchiyama et al. 2012a, b).

*Other Effects on Animal Organisms*

Several effects on animal behavior in different Vertebrates have been reviewed (Dhungana et al. 2018). Studies in chicken have shown that music reduces stress measured by various parameters (duration of tonic immobility, white blood cells/lymphocyte ratio and fluctuating asymmetry of the organism) (Bonzom 1999, Dávila et al. 2011). It is interesting to observe that the effects described according to the type of music and the animal species can be positive, negative or inexistent. In birds, it is known that the young learn the characteristic melodies of their species by listening to the adults. Young birds deprived of sound, either because they are deaf or because they are raised in isolation, away from the song of adults, are unable to sing. The learning of birdsong in relation to the intrinsic plasticity of neurons has been studied recently in zebra finch (Daou and Margoliash 2021).

Several results were also been obtained in monkeys: music can have effects in decreasing the heartrate of baboons, increasing the social behaviour in chimpanzees, or inducing an abnormal behaviour in rhesus macaques (Dhungana et al. 2018). In rats, music might improve spatial learning and memory ability (Xiong et al. 2018).

Effects of music have also been observed in domestic animals. In dogs, classical music can decrease vocalization and body shaking and increase sleep duration. Cats are sensitive to music and suitable musical pieces have been created (Dhungana et al. 2018). On the other hand, various effects have been observed in cattle. It has been observed that dairy breeds are more sensitive to sound, and especially music, than beef breeds (Dhungana et al. 2018). Increase or decrease in milk production depends on the noise levels. Loud noise or excessive sounds cause a stress of cattle and therefore a decrease in the amount of milk produced (Hemsworth 2003). In contrast, slow music can increase milk production contrarily to fast music (Algers et al. 1978, Algers and Jensen 1991). In buffalo, animal behaviour and reproduction are affected by music (Abuzead and Khalil 2007).

On the other hand, music seems to have effects on embryonic development, facilitating neurogenesis, and also regeneration and repair of neurons in Human (Kim et al. 2006, Fukui and Toyoshima 2008), and modulating expression of apoptosis in the development of auditory nuclei in chicks (Alladi et al. 2005). In a study devoted to the effects of music on the development, pairs of Wistar rats were exposed to classic music. Before parturition, the fetuses were extracted and brains prepared for electron microscopy. The results showed that prenatal music had an effect on neuroplasticity of fetal brain. As rat fetuses cannot hear until birth, these neuroplasticity changes of fetal brain could be related to a reduction in the mothers' blood corticosterone level (Sheikhi and Saboory 2015). In chicks, prenatal music stimulation leads to morphological changes in hippocampus and brainstem auditory nuclei. In contrast, unpatterned noise and loud music (110 dB) induce altered volume and neuronal number in these regions, indicating that the characteristic of the sound was mediating effects on brain development (Sanyal et al. 2013). Furthermore, it has been shown that *in ovo* sound stimulation (music or noise: 110 dB) affects the regulation of brain-derived neurotrophic factor (BDNF) in the

developing auditory system of chicks; it was regulated at the transcriptional and post-translational level in the auditory cortex and at the post-translational level in the hippocampus (Kathpalia et al. 2019).

According to some authors, music would also improve the success of fertilization rates of *in vitro* cultured human embryos (Lopez-Teijón et al. 2015).

At last, some studies have shown the effects of sounds and music on stem cells (Choi et al. 2016, Wei et al. 2016, Ventura et al. 2017).

#### *Some Adverse Effects on Animal Cells*

The effects of music cannot be equated with the effects of all types of sound. Research shows that not all types of sound have beneficial effects on the cells. There are sounds with harmful effects (Clark 1991). When the choroidal plexus of rats is submitted to noise, the number of normal cells decreased and the number of apoptotic cells increased (McCarthy et al. 1992, Aydin et al. 2011). In experiments conducted to analyze the effects of anxiolytics in rats, both noise and music suppressed the sexual behavior of females. In another experiment, noise suppressed the sexual impulses of ovariectomized females treated with high dose of estradiol and induced avoidance of the area where males were located (Le Moëne and Ågmo 2018a, b, Le Moëne et al. 2019). These studies show that in some cases, music can negatively affect animal behavior.

#### *Effects on Unicellular Organisms*

Like multicellular organisms, unicellular organisms, eukaryotes or prokaryotes, seem to be sensitive to music. Several works have shown the effects of single-frequency sound on the biology of various bacteria (Mortazavian 2012). Experiments using indian music have shown that it has a positive effect on the growth of bacteria and yeasts. Beneficial effects have thus been shown on cellular growth, metabolism and antibiotic sensitivity (Norris and Hyland 1997, Matsushashi et al. 1998, Ayan et al. 2008, Shaobin et al. 2010, Sarvaiya and Kothari 2015, Shah et al. 2016). Growth, pigment production and antibiotic sensibility are affected by the intensity of sound in several bacteria (Kothari and Sarvaiya 2017, Khotari et al. 2018).

#### *Effects on Plants*

In plants, sound waves can influence germination, development and growth (Qin et al. 2003, Mishra et al. 2016, Vicient 2017). It can also enhance the immunity increasing tolerance to drought (Chivukula and Ramaswamy 2014, Choi et al. 2017, Lopez-Ribera and Vicient 2017). A recent study looked at the response of *Oenothera drummondii* plants to sound. The main result obtained is that the flowers of this plant produce a sweeter nectar only three minutes after being exposed to the rustling of the wings of butterflies and bees which are its pollinators (De Luca and Vallejo-Marin 2013, Veits et al. 2019). Other examples are consistent with the existence of an acoustic communication during interactions between



plants and pollinators (Ayan et al. 2008, Mishra et al. 2016, Schöner et al. 2016). More broadly, some animals and other plants can use the sounds emitted by a plant to obtain information about its condition (Khait et al. 2019).

Some studies have shown that the roots of various plants orient themselves towards the noise generated for example by flowing water (Gagliano et al. 2012, Gagliano 2013) and that the corn roots tend to grow towards a sound source whose frequency is close to 200 Hz (Gagliano 2013). Those different examples confirm that plants can also use sound, but the ecological and evolutionary implications that this represents in the life of a plant are not yet well known (Hongbo et al. 2008, Gagliano et al. 2012, Gagliano 2013). Further studies on plant bioacoustics are still needed to understand the interactions of plants with their environment.

### *Effects on Genome*

Studies have looked at the roles of various forms of genomic changes and gene regulation related to music abilities. Some genes inherited with musical aptitude have been identified. In particular, the alpha-synuclein gene, involved in song perception in songbirds, has been shown to be also overexpressed in individuals listening or performing music. Its mutation is associated to several cognitive diseases in humans and perturbations of song production and perception in songbirds. These observations are in accordance with an evolutionary conservation in biological processes related to sound perception (Kanduri et al. 2015, Järvellä 2018). On the other hand, changes in the peripheral blood microRNA transcriptome after 2 hours of concert performance are identified in professional musicians, suggesting regulatory mechanisms underlying auditory perception. Based on the current knowledge about the microRNA function in blood and brain, the authors interpreted the regulation of these non-coding RNAs as consequence of a regular musical activity. In brain, the putative targets of these microRNAs were involved in neuronal activity, neuronal plasticity, myelination and cognitive functions (Nair et al. 2019, Nair et al. 2021).

It has been shown that Indian Vedic music can upregulate the expression of steroidogenic genes in 3D cultured buffalo granulosa cell spheroids (Pandey et al. 2021).

Proliferation of gastric cancer cells did not increase when cells were exposed to classical music contrarily to those exposed to metal music. The expression of several genes involved in proliferation seems to be modulated in different ways according to the music type (Ramírez-Rivera and Bernal 2019).

In order to give a molecular basis to effects of music on organisms, a meta-analysis of 105 publications concerning humans and several animal models identified several candidate genes related to musical traits. The authors have shown that these genes are involved in biological functions underlying learning and memory, which are essential properties for music abilities (Oikkonen et al. 2016). A study of properties of audible sound as mechanical stimulus by cells has shown an effect of wave form and sound pressure level, more than frequency, on gene expression, with a cell type-specific response to sound (Kumeta et al. 2018).

## **Reception of Vibrations and Intracellular Repercussions**

How vibrations can intervene on cellular physiology? The effects of vibrations are well known in hearing cells. These cells also called “hair cells” are sensory cells emitting cilia which are really cytoplasmic expansions, the stereocilia. These cells are arranged along the basilar membrane of the cochlea (still called “organ of Corti”). Depending on their position on the cochlea, cell membranes are sensitive to vibrations at well-defined frequencies. The vibrations cause a deformation of the membrane of these cells at the level of the cilia which amplifies the surface of reception of the vibrations. The deviation of the stereocilia causes the opening of ion channels creating an electrical receptor potential.  $\text{Ca}^{++}$  ions enter the cell, releasing glutamate, a neurotransmitter at the level of the synapses in connection with axons of neurons. The signal is transmitted to the auditory areas of the brain. This type of interactions between cell membrane and vibrations can certainly be generalized to all cells.

All cell membranes are made up of a bilayer of phospholipids in which proteins are incorporated. This membrane is bound outside the cell and can carry receptors; it is also linked directly to the cytoskeleton and to the enzyme systems located inside the cell. Even if studies specifically related to the mechanism of the effects of the sounds received by cell membranes are still lacking, the analysis of works on membrane receptors, the nature of the signals and the elastic capacities of the cell allow us to make assumptions as to these mechanisms.

At the cellular level, it is certainly in plants that work is the most advanced. Sound vibrations can affect microfilament rearrangements, increase the levels of polyamines and soluble sugars, alter the activity of various proteins, and regulate the transcription of some genes (Qin et al. 2003, Hongbo et al. 2008, Mishra et al. 2016). The reception of chemical signals by animal cells is well known. Various stimuli such as light, ions, pheromones, hormones or neurotransmitters, bind to G protein-coupled receptors located on cell membranes. Then, they transduce these extracellular signals inside the cells by engaging G proteins, that trigger a cascade of signaling events leading to an appropriate cellular response (Lim et al. 2015, Syrovatkina et al. 2016). Further studies on the modalities of acoustic signal capture will provide a better understanding of the intracellular repercussions of sound. Elasticity of biological tissues can be measured by elastography. This imaging method makes it possible to understand the variations in the hardness of tissues and their elasticity (Brum 2012, Grasland-Mongrain 2013) as a consequence of the links that membrane receptors establish with the elements of the cytoskeleton (Icard-Arcizet 2007).

Reading these works, we can assume that sound vibrations trigger the plasmatic membrane’s vibration, which could result in the activation of certain proteins of the G protein type and the transmission of a chemical signal in the cell. Several metabolic reactions governing the physiology of the cell could then be affected.

A controversial theory, the “proteody theory”, emitted by the French particle physicist Joël Sternheimer at the beginning of the 1990s, defended the idea of the existence of waves emitted in the body molecules. Unfortunately, no scientific

publications are available for now (Doux 2019). According to this theory, waves whose frequencies can be transcribed into musical notes are associated with each group of amino acids. In the same vein, King and Angus (1996) published a computer program called PM (protein music) to analyze information about protein sequences by audification, i.e. using the hearing to analyze data (Kramer 1994).

Petrovski et al. (2021) found a correlation between yeast vacuolar proton-ATPase activity (involved in many life processes) and the intensity of several frequencies of an oscillating electric field generated by music. These observations led these authors to conclude that such an electric field from the music interfered with the periodic transmembrane movements of the enzyme. Based on the fact that organisms are subject in nature to several rhythmic oscillations, Muehsam and Ventura (2014) give a theoretical study in which they examine the effects of sound vibrations on gene expression.

## Conclusions

During biological evolution, organisms have acquired senses that allow them to interact with the environment. The sensory perception of living organisms is very diversified and is directly related to the lifestyle of the organism. Several of those senses will be put to good use by art in Human. Among the arts, music is directly related to the sensory system of hearing. While the effects of sound vibrations on auditory cells have long been known, the effects on other cell types are much less well understood. Currently, an increasing number of works is investigating the effects of music on the biochemistry of non-auditory cells and on various aspects of metabolism. Nevertheless, these works remain still quite disparate.

Music elicits reactions in humans that promote positive emotions, stress relief and immune function, by soliciting different cellular signaling molecules including hormones, neurotransmitters, cytokines, and peptides (Gangrade 2012). Music can restore some of the homeostasis and thus reduce pain (Nelson et al. 2008). It probably reduces alterations in the hypothalamic-pituitary hormonal axes (Russo et al. 2017). Music (432 Hz) can also influence growth, by increasing food intake and body weight; a significant increase of ghrelin expression was found in hypothalamic neurons in music-treated rat pups (Russo et al. 2021). A hypothesis involves quantum mechanics. But beware of the enthusiasm aroused by these exciting results. Studies are still necessary to understand and go further in the possible applications, the precise molecular mechanisms of its effects indeed yet to be clarified. But there is a field of research emerging here.

## References

- Abuzead SMM, Khalil AM (2007) Behavioral and physiological influences of listening slow and fast music on milking buffaloes. *Assiut Veterinary Medicine Journal* 53(114): 1–13.
- Albanese A, Battisti E, Vannoni D, Acetoi E, Galassi G, Giglioni S, et al. (2009) Alterations in adenylate kinase activity in human PBMCs after *in vitro* exposure to

- electromagnetic field: comparison between extremely low frequency electromagnetic field (ELF) and therapeutic application of a musically modulated electromagnetic field (TAMMEF). *Journal of Biomedicine and Biotechnology* 2009(Sep): 717941.
- Algers B, Jensen P (1991) Teat stimulation and milk production during early lactation in sows: effects of continuous noise. *Canadian Journal of Animal Science* 71(1): 51–60.
- Algers B, Ekesbo I, Strömberg S (1978) The impact of continuous noise on animal health. *Acta Veterinaria Scandinavica. Supplementum* 1978(68): 1–26.
- Alladi PA, Roy T, Singh N, Wadhwa S (2005) Prenatal auditory enrichment with species specific calls and sitar music modulates expression of *Bcl-2* and *Bax* to alter programmed cell death in developing chick auditory nuclei. *International Journal of Developmental Neurosciences* 23(4): 363–373.
- Arai MN (1997) *A functional biology of Scyphozoa*. London, UK: Chapman and Hall.
- Ashmore J (2008) Cochlear outer hair cell motility. *Physiological Review* 88(1): 173–210.
- Ayan I, Aslan G, Cömelekoğlu U, Yilmaz N, Colak M (2008) The effect of low intensity pulsed sound waves delivered by the exogen device on *Staphylococcus aureus* morphology and genetics. (In Turkish). *Acta Orthopaedica et Traumatologica Turcica* 42(4): 272–277.
- Aydin MU, Ungoren MK, Aydin N, Halici Z, Onder A, Gundogdu C, et al. (2011) The effects of impulse noise on the epithelial cells of the choroid plexus. *Turkish Neurosurgery* 21(2): 191–196.
- Badenhorst A (1978) The development and the phylogeny of the organ of Jacobson and the tentacular apparatus of *Ichtyophis glutinosus* (Linn.) *Annals of the University of Stellenbosch* 1(1): 1–26.
- Barnes RSK, Calow PC, Olive PJW, Golding DW, Spicer JI (2009) *The invertebrates—A synthesis*. 3rd Edition. Oxford, UK: Wiley.
- Behzadmehr R, Dastyar N, Moghadam MP, Abavisani M, Moradi M (2020) Effect of complementary and alternative medicine interventions on cancer related pain among breast cancer patients: a systematic review. *Complementary Therapies in Medicine* 49(Mar): 102318.
- Billar RJ, Kühlmann AYR, Schnater JM, Vlot J, Tomas JJP, Zijp GW, et al. (2020) Interventions with music in pectus excavatum treatment (IMPECT trial): a study protocol for a randomised controlled trial investigating the clinical effects of perioperative music interventions. *BMJ Open* 10(7): e036380.
- Billo R (1986) The tentacle apparatus of Caecilians. *Mémoires de la Société Zoologique de France* 43: 71–75.
- Billo R, Wake MH (1987) Tentacle development of *Dermophis mexicanus* (Amphibian: Gymnophiona) with an hypothesis on tentacle origin. *Journal of Morphology* 192(2): 101–111.
- Bittman BB, Berk LS, Felten DL, Westengard J, Simonton OC Pappas J, et al. (2001) Composite effects of group drumming music therapy on modulation of neuroendocrine-immune parameters in normal subjects. *Alternative Therapies in Health and Medicine* 7(1): 38–47.
- Bleckmann H, Zelick R (2009) Lateral line system of fish. *Integrative Zoology* 4(1): 13–25.
- Bolgan M, Gervaise C, di Iorio L, Lossent J, Lejeune P, Raick X, et al. (2020) Fish biophony in a Mediterranean submarine canyon. *The Journal of the Acoustical Society of America* 147(4): 2466.
- Bonzom JM (1999) *L'asymétrie fluctuante un biomarqueur morphométrique pour évaluer la qualité de l'environnement*. (Fluctuating asymmetry a morphometric biomarker to assess environmental quality). PhD Thesis. Canada: University of Sherbrook.

- Bowman A, Scottish SPCA, Dowell FJ, Evans NP (2017) The effect of different genres of music on the stress levels of kennelled dogs. *Physiology & Behavior* 15(171): 207–215.
- Brum J (2012) *Elastographie et retournement temporel des ondes de cisaillement : application à l'imagerie des solides mous*. (Elastography and time reversal of shear waves: application to the imaging of soft solids). PhD Thesis. France: Université de Grenoble.
- Buglione A, Saccone G, Mas M, Raffone A, di Meglio L, di Meglio L, et al. (2020) Effect of music on labor and delivery in nulliparous singleton pregnancies: a randomized clinical trial. *Archives of Gynecology and Obstetrics* 301(3): 693–698.
- Bulut M, Alemdar DK, Bulut A, Şalcı G (2020) The effect of music therapy, hand massage, and kaleidoscope usage on postoperative nausea and vomiting, pain, fear, and stress in children: a randomized controlled trial. *Journal of Perianesthesia Nursing* 35(6): 649–657.
- Cardinale M, Pope MH (2003) The effects of whole body vibration on humans: dangerous or advantageous? *Acta Physiologica Hungarica* 90(3): 195–206.
- Çelebi D, Yılmaz E, Şahin ST, Baydur H (2020) The effect of music therapy during colonoscopy on pain, anxiety and patient comfort: a randomized controlled trial. *Complementary Therapies in Clinical Practices* 2020(38): 101084.
- Chai PR, Gale JY, Patton ME, Schwartz E, Jambaulikar GD, Taylor SW, et al. (2020a) The impact of music on nociceptive processing. *Pain Medicine* 21(11): 3047–3054.
- Chai PR, Schwartz E, Hasdianda MA, Azizoddin DR, Kikut A, Jambaulikar GD, et al. (2020b) A brief music app to address pain in the emergency department: prospective study. *Journal of Medical Internet Research* 22(5): e18537.
- Chan DK, Hudspeth AJ (2005)  $Ca^{2+}$  current driven nonlinear amplification by the mammalian cochlea *in vitro*. *Nature Neuroscience* 8(2): 149–155.
- Chan JCH, Chan LP, Yeung CP, Tang TW, O YM, Lam, WC (2020) Effect of music on patient experience during intravitreal injection. *Journal of Ophthalmology* 2020(13): 1–6.
- Chivukula V, Ramaswamy S (2014) Effect of different types of music on *Rosa chinensis* plants. *International Journal of Environmental Science and Development* 5(5): 431–434.
- Choi B, Ghosh R, Gururani MA, Shanmugam G, Jeon J, Kim J, et al. (2017) Positive regulatory role of sound vibration treatment in *Arabidopsis thaliana* against *Botrytis cinerea* infection. *Science Reports* 7: 2527.
- Choi Y, Park JE, Jeong JS, Park JK, Kim J, Jeon S (2016) Sound waves induce neural differentiation of human bone marrow-derived mesenchymal stem cells *via* Ryanodine receptor-induced calcium release and Pyk2 activation. *Applied Biochemistry and Biotechnology* 180(4): 682–694.
- Cimen SG, Oğuz E, Gundogmus AG, Cimen S, Sandikci F, Ayli MD (2020) Listening to music during arteriovenous fistula surgery alleviates anxiety: a randomized single-blind clinical trial. *World Journal of Transplantation* 10(4): 79–89.
- Clark WW (1991) Noise exposure from leisure activities: a review. *The Journal of the Acoustical Society of America* 90(1): 175–181.
- Corallo C, Volpi N, Franci D, Vannoni D, Leoncini R, Landi G, et al. (2013) Human osteoarthritic chondrocytes exposed to extremely low-frequency electromagnetic fields (ELF) and therapeutic application of musically modulated electromagnetic fields (TAMMEF) systems: a comparative study. *Rheumatology International* 33(6): 1567–1575.
- Corallo C, Battisti E, Albanese A, Vannoni D, Leoncini R, Landi G, et al. (2014) Proteomics of human primary osteoarthritic chondrocytes exposed to extremely low-

- frequency electromagnetic fields (ELF EMFs) and to therapeutic application of musically modulated electromagnetic fields (TAMMEF). *Electromagnetic Biology and Medicine* 33(1): 3–10.
- Dai WS, Huang ST, Xu N, Chen Q, Cao H (2020) The effect of music therapy on pain, anxiety and depression in patients after coronary artery bypass grafting. *Journal of Cardiothoracic Surgery* 15(1): 81.
- Daou A, Margoliash D (2021) Intrinsic plasticity and birdsong learning. *Neurobiology of Learning and Memory* 180(Apr): 107407.
- Dávila SG, Campo JL, Gil MG, Prieto MT, Torres O (2011) Effects of auditory and physical enrichment on 3 measurements of fear and stress (tonic immobility duration, heterophil to lymphocyte ratio, and fluctuating asymmetry) in several breeds of layer chicks. *Poultry Science* 90(11): 2459–2466.
- De Luca PA, Vallejo-Marin M (2013) What's the 'buzz' about? The ecology and evolutionary significance of buzz pollination. *Current Opinion in Plant Biology* 16(4): 429–435.
- Dhungana S, Khanal DR, Sharma M, Bhattarai N, Tamang DT, Wasti S, et al. (2018) Effect of music on animal behavior: a review. *Nepalese Veterinary Journal* 35(Aug): 142–149.
- Di Iorio L, Dailianis T, Catalan I, Lejeune P, Parmentier E (2019) An acoustic odyssey: characterisation of the vocal fish community inhabiting Neptune seagrass meadows across the Mediterranean Sea. In *International Bioacoustics Congress*. Brighton, UK.
- Doux A (2019, April 11) *Les protéodies, la musique du vivant*. (Proteodies, the music of life). Des musiques pour guérir. Retrieved from: <https://desmusiquespourguerir.com/les-proteodies-la-musique-du-vivant/>. [Accessed 5 June 2021]
- Døving KB, Trotier D (1998) Structure and function of the vomeronasal organ. *Journal of Experimental Biology* 201(21): 2913–2925.
- Dunbar RIM, Pearce E, Tarr B, Makdani A, Bamford J, Smith S, et al. (2021) Cochlear SGN neurons elevate pain thresholds in response to music. *Scientific Reports* 11(1): 14547.
- Erken G, Bor Kucukatay M, Erken HA, Kursunluoglu R, Genc O (2008) Influence of classical and rock music on red blood cell rheological properties in rats. *Medical Science Monitor* 14(1): 28–33.
- Ernberg M, Al-Khdhairi D, Shkola K, Jounger SL, Christidis N (2020) Experimental muscle pain and music, do they interact? *Journal of Oral Pathology and Medicine* 49(6): 522–528.
- Exbrayat JM, Brun C (2019) Some effects of sound and music on organisms and cells: a review. *Annual Research and Review in Biology* 32(2): 1–12.
- Fettiplace R, Hackney CM (2006) The sensory and motor roles of auditory hair cells. *Nature Neuroscience* 7(1): 19–29.
- Fish MP, Mowbray WH (1970) *Sounds of western North Atlantic fishes; a reference file of biological underwater sounds*. Baltimore, Maryland: Johns Hopkins Press.
- Fukui H, Toyoshima AK (2008) Music facilitates the neurogenesis, regeneration and repair of neurons. *Medical Hypotheses* 71(5): 765–769.
- Gagliano M (2013) Green symphonies: a call for studies on acoustic communication in plants. *Behavioral Ecology* 24(4): 789–796.
- Gagliano M, Renton M, Duvdevani N, Timmins M, Mancuso S (2012) Acoustic and magnetic communication in plants: Is it possible? *Plant Signaling and Behavior* 7(10): 1346–1348.
- Gamboa J, Cameron MC, Fathi R, Alkousakis T (2020) A review of non-pharmacologic approaches to enhance the patient experience in dermatologic surgery. *Dermatology Online Journal* 26(3): 13030/qt7mp372nd.

- Gangrade A (2012) The effect of music on the production of neurotransmitters, hormones, cytokines, and peptides: a review. *Music and Medicine* 4(1): 40–43.
- Garcia Guerra G, Joffe A, Sheppard C, Hewson K, Dinu IA, de Caen A, et al. (2020) Canadian Critical Care Trials Group. Music use for sedation in critically ill children (MUSiCC trial): study protocol for a pilot randomized controlled trial. *Pilot and Feasibility Studies* 6: 31.
- Gibbs MA, Northcutt RG (2004) Development of the lateral line system in the shovelnose sturgeon. *Brain, Behavior and Evolution* 64(2): 70–84.
- Giordano F, Scarlata E, Baroni M, Gentile E, Puntillo F, Brienza N, et al. (2020) Receptive music therapy to reduce stress and improve wellbeing in Italian clinical staff involved in COVID-19 pandemic: a preliminary study. *The Arts in Psychotherapy* 70(Jul): 101688.
- Gogoularadja A, Bakshi SS (2020) A randomized study on the efficacy of music therapy on pain and anxiety in nasal septal surgery. *International Archives of Otorhinolaryngology* 24(2): e232–e236.
- González-Grajales LA, Pieper L, Görner P, Görner S, Staufenbiel R (2019) Effects of auditory and visual stimuli on glucose metabolism in Holstein dairy cattle. *Acta Veterinaria Scandinavica* 61(1): 2.
- Grasland-Mongrain P (2013) *Applications de la force de Lorentz en acoustique médicale*. (Applications of the Lorentz force in medical acoustics). PhD Thesis. France: University Lyon 1.
- Guo S, Lu J, Wang Y, Li Y, Huang B, Zhang Y, et al. (2020) Sad music modulates pain perception: an EEG study. *Journal of Pain Research* 13(Aug): 2003–2012.
- Hallmann A (2003) Extracellular matrix and sex-inducing pheromone in *Volvox*. *International Review of Cytology* 227(Feb): 131–182.
- Hasegawa Y, Kubota N, Inagaki T, Shinagawa N (2001) Music therapy induced alternations in natural killer cell count and function. (In Japanese). *Nihon Ronen Igakkai Zasshi* 38(2): 201–204.
- Hemsworth PH (2003) Human–animal interactions in livestock production. *Applied Animal Behaviour Science* 81(3): 185–198.
- Hirokawa E, Ohira H (2003) The effects of music listening after a stressful task on immune functions, neuroendocrine responses, and emotional states in college students. *Journal of Music Therapy* 40(3): 189–211.
- Hongbo S, Biao L, Bochu W, Kun T, Yilong L (2008) A study on differentially expressed gene screening of Chrysanthemum plants under sound stress. *Comptes Rendus Biologies* 331(5): 329–333.
- Howlin C, Rooney B (2020) The cognitive mechanisms in music listening interventions for pain: a scoping review. *Journal of Music Therapy* 57(2): 127–167.
- Huby A, Lowie A, Herrel A, Vigouroux R, Frederic B, Raick X, et al. (2019) Functional diversity in biters: the evolutionary morphology of the oral jaw system in pacus, piranhas and relatives (Teleostei: Serrasalminidae). *Biological Journal of the Linnean Society* 127(4): 722–741.
- Hurnik JF, Johnson L (1997) Effect of music on voluntary approach of dairy cows to an automatic milking system. *Applied Animal Behaviour Science* 53(3): 175–182.
- Icard-Arcizet D (2007) *Modifications mécaniques et biologiques induites dans des cellules en culture par l'application locale d'une force contrôlée*. (Mechanical and biological changes induced in cells in culture by the local application of a controlled force). PhD Thesis. France: Université Paris-Diderot.
- Ikushima N, Maruyama S (2007) The protoplasmic connection in *Volvox*. *Journal of Eukaryotic Microbiology* 15(1): 136–140.

- Järvelä I (2018) Genomics studies on musical aptitude, music perception, and practice. *Annals of the New York Academy of Sciences* 1423(1): 82–91.
- Johnson AJ, Elkins GR (2020) Effects of music and relaxation suggestions on experimental pain. *International Journal of Clinical and Experimental Hypnosis* 68(2): 225–245.
- Jones WD, Kalantari A, Otte A (2020) Neurostimulation durch Musik. (Neurostimulation through music). *MMW Fortschritte der Medizin* 162(suppl 4): 3–8.
- Kanduri C, Kuusi T, Ahvenainen M, Philips AK, Lähdesmäki H, Järvelä I (2015) The effect of music performance on the transcriptome of professional musicians. *Scientific Reports* 25(5): 9506.
- Kathpalia P, Nag TN, Chattopadhyay P, Sharma A, Bhat MA, Roy TS, et al. (2019) *In ovo* sound stimulation mediated regulation of BDNF in the auditory cortex and hippocampus of neonatal chicks. *Neuroscience* 408(Apr): 293–307.
- Kemp A (2020) *The effects of music on dairy production*. PhD Thesis. Kentucky, USA: Murray State University.
- Kenison L (2016) *The effects of classical music on dairy cattle*. PhD Thesis. USA: Alfred University.
- Khait I, Lewin-Epstein O, Sharon R, Saban K, Perelman R, Boonman A, et al. (2019) *Plants emit informative airborne sounds under stress*. Preprint BioRxiv 2019.
- Kim H, Lee MH, Chang HK, Lee TH, Lee HH, Shin MC, et al. (2006) Influence of prenatal noise and music on the spatial memory and neurogenesis in the hippocampus of developing rats. *Brain and Development* 28(Mar): 109–114.
- King RD, Angus CG (1996) PM - Protein music. *Cabios Applications Note* 12(3): 251–252.
- King B, Hu H, Long JA (2018) Electroreception in early vertebrates: survey, evidence and new information. *Palaeontology* 61(3): 325–358.
- Köhler F, Martin ZF, Hertrampf RS, Gäbel C, Kessler J, Ditzen B, et al. (2020) Music therapy in the psychosocial treatment of adult cancer patients: a systematic review and meta-analysis. *Frontiers in Psychology* 11(Apr): 651.
- Kothari V, Sarvaiya N (2017) Audible sound in form of music can influence microbial growth, metabolism and antibiotic susceptibility. *Journal of Applied Biotechnology and Bioengineering* 2(19): 212–219.
- Kothari V, Joshi C, Patel P, Mehta M, Dubey S, Mishra B, et al. (2018) Influence of a Mono-Frequency Sound on Bacteria can be a Function of the Sound-Level. *Indian Journal of Science and Technology* 11(4): 1–9.
- Kramer G (1994) *Auditory display: sonification, audification, and auditory interface*. Boca Raton, USA, London, UK, New York, USA: CRC Press.
- Kumeta M, Takahashi D, Takeyasu K, Yoshimura SH (2018) Cell type-specific suppression of mechanosensitive genes by audible sound stimulation. *PLoS One* 13(1): e0188764.
- Le Moëne O, Ågmo A (2018a) Responses to positive and aversive stimuli in estrous female rats housed in a seminatural environment: Effects of yohimbine and chlordiazepoxide. *Pharmacology, Biochemistry and Behavior* 179(Apr): 43–54.
- Le Moëne O, Ågmo A (2018b) Behavioral responses to emotional challenges in female rats living in a seminatural environment: the role of estrogen receptors. *Hormones and Behavior* 106(Nov): 162–177.
- Le Moëne O, Stavarache M, Ogawa S, Musatov S, Ågmo A (2019) Estrogen receptors  $\alpha$  and  $\beta$  in the central amygdala and the ventromedial nucleus of the hypothalamus: sociosexual behaviors, fear and arousal in female rats during emotionally challenging events. *Behavioural Brain Research* 367(Jul): 128–142.
- Lemarquis P (2009) *Sérénade pour un cerveau musicien*. (Serenade for a brain musician). Paris, France: Odile Jacob.



- Lemarquis P (2021) *Les pouvoirs de la musique sur le cerveau des enfants et des adultes*. (The powers of music on the brain of children and adults). Paris, France: Odile Jacob.
- Lestard NR, Capella MA (2016) Exposure to music alters cell viability and cell motility of human non-auditory cells in culture. *Evidence-Based Complementary and Alternative Medicine* 2016(23): 1–7.
- Lestard NR, Valente RC, Lopes AG, Capella MA (2013) Direct effects of music in non-auditory cells in culture. *Noise Health* 15(66): 307–314.
- Lim W, Mayer B, Pawson T (2015) *Cell signaling: principles and mechanisms*. New York, US: Garland Science.
- Lopez-Ribera I, Vicient CM (2017) Drought tolerance induced by sound in Arabidopsis plants. *Plant Signaling and Behavior* 12(10): e1368938.
- Lopez-Teijón M, Castelló C, Asencio M, Fernández P, Farreras A, Rovira S, et al. (2015) Improvement of fertilization rates of in vitro cultured human embryos by exposure to sound vibrations. *Journal of Fertilization: In Vitro-IVF-Worldwide, Reproductive Medicine, Genetics and Stem Cell Biology* 3(4): 1000160.
- Lu Y, Liu M, Shi S, Jiang H, Yang L, Liu X, et al. (2010) Effects of stress in early life on immune functions in rats with asthma and the effects of music therapy. *Journal of Asthma* 47(4): 526–531.
- Lu SY, Peng GY, Gu LG, Li ZM, Yin SJ (2013) Effect and mechanisms of gong-tone music on the immunological function in rats with Liver (Gan)-qi depression and Spleen (Pi)-qi deficiency syndrome in rats. *Chinese Journal of Integrative Medicine* 19(3): 212–216.
- Martin J, Le Faucheur L, Pondarré C, Carlier-Gonod A, Baeza-Velasco C (2020) Effets d'une intervention musicale standardisée sur la gestion de la douleur et de l'anxiété-état des adolescents drépanocytaires. (Effects of a standardized music intervention on the management of pain and state-anxiety in adolescents with sickle cell disease). *Soins Pédiatrie/Puericulture* 41(314): 35–41.
- Matsushashi M, Pankrushina AN, Takeuchi S, Ohshima H, Miyoi H, Endoh K, et al. (1998) Production of sound waves by bacterial cells and the response of bacterial cells to sound. *The Journal of General and Applied Microbiology* 44(1): 49–55.
- Mayoud L, Lemarquis P (2019) *L'invitation à la beauté: L'ouverture au monde par l'empathie esthétique*. (The invitation to beauty: openness to the world through aesthetic empathy). Paris et Lyon, France: Vrin et IIEE.
- McCarthy DO, Ouimet ME, Daun JM (1992) The effects of noise stress on leukocyte function in rats. *Research in Nursing and Health* 15(2): 131–137.
- Mehr SA, Krasnow MM, Bryant GA, Hagen EH (2020) Origins of music in credible signaling. *Behavioral and Brain Sciences* 44(Aug): e60.
- Mishra RC, Ghosh R, Bae H (2016) Plant acoustics: in the search of a sound mechanism for sound signaling in plants. *Journal of Experimental Botany* 67(15): 4483–4494.
- Mohammed T, Murphy MF, Lilley F, Burton DR, Bezombes F (2016) The effects of acoustic vibration on fibroblast cell migration. *Materials Science and Engineering C: Materials for Biological Applications* 69(Dec): 1256–1262.
- Moore J (2006) *An introduction to the invertebrates*. Cambridge, UK: Cambridge University Press.
- Mortazavian AM (2012) Music affects survival and activity of microorganisms. *Archives of Advances in Biosciences* 3(4): 1.
- Moss CF, Sinha SR (2003) Neurobiology of echolocation in bats. *Current Opinion in Neurobiology* 13(6): 751–758.
- Muehsam D, Ventura C (2014) Life rhythm as a symphony of oscillatory patterns: electromagnetic energy and sound vibration modulates gene expression for biological signaling and healing. *Global Advances in Health and Medicine* 3(2): 40–55.

- Müller U (2008) Cadherins and mechanotransduction by hair cells. *Current Opinion in Cell Biology* 20(5): 557–566.
- Nair PS, Kuusi T, Ahvenainen M, Philips AK, Järvelä I (2019) Music-performance regulates microRNAs in professional musicians. *PeerJ* 7(1): e6660.
- Nair PS, Rajjas P, Ahvenainen M, Philips AK, Ukkola-Vuoti L, Irma Järvelä I (2021) Music-listening regulates human microRNA expression. *Epigenetics* 16(5): 554–566.
- National Research Council (US) Committee on Assessment of the Possible Health Effects of Ground Wave Emergency Network (1993) *Assessment of the possible health effects of ground wave emergency network*. Washington DC, US: National Academies Press.
- Nelson A, Hartl W, Jauch KW, Fricchione GL, Benson H, Warshaw AL, et al. (2008) The impact of music on hypermetabolism in critical illness. *Current Opinion in Clinical Nutrition and Metabolic Care* 11(6): 790–794.
- Norris V, Hyland GJ (1997) Do bacteria sing? Sonic intercellular communication between bacteria may reflect electromagnetic intracellular communication involving coherent collective vibrational modes that could integrate enzyme activities and gene expression. *Molecular Microbiology* 24(4): 879–880.
- Núñez MJ, Maña P, Liñares D, Riveiro MP, Balboa J, Suarez-Quintanilla J, et al. (2002) Music, immunity and cancer. *Life Sciences* 71(9): 1047–1057.
- Oikkonen J, Onkamo P, Järvelä I, Kanduri C (2016) Convergent evidence for the molecular basis of musical traits. *Scientific Reports* 6(1): 39707.
- Pandey M, Singh C, Goud ESK, Vadamurthy VG, Singh D, Onteru SK (2021) Effect of Vedic music on steroidogenic gene expression in 3D-cultured buffalo granulosa cell spheroids model system, a pilot study. *Reproduction in Domestic Animals* 56(2): 231–238.
- Parmentier E, Lecchini D, Mann D (2016) Sound production in damselfishes. In E Parmentier, B Frédérick (eds.), *Biology of Damselfishes*, 204–228. New York, US: CRC Press.
- Payne R (1984) *Communication and behavior of whales*. Boulder, Col, US: Westview Press.
- Pérez-Eizaguirre, M, Vergara-Moragues E (2020) Music therapy interventions in palliative care: A systematic review. *Journal of Palliative Care* 36(3): 194–205.
- Petrovszki P, Sebök-Nagy K, Páli T (2021) The activity of native vacuolar proton-ATPase in an oscillating electric field - Demystifying an apparent effect of music on a biomolecule. *Frontiers in Molecular Biosciences* 8(Nov): 772167.
- Polascik BA, Tan DJA, Raghunathan K, Kee HM, Lee A, Sng BL, et al. (2021) Acceptability and feasibility of perioperative music listening: a rapid qualitative inquiry approach. *Journal of Music Therapy* 58(1): 43–69.
- Pulcrano S, di Porzio U (2021) The alchemy of music and the brain. *Academia Letters*, Art 4185.
- Qin YC, Lee WC, Choi YC, Kim TW (2003) Biochemical and physiological changes in plants as a result of different sonic exposures. *Ultrasonics* 41(5): 407–411.
- Rabbitt RD, Boyle R, Highstein SM (2010) Mechanical amplification by hair cells in the semicircular canals. *PNAS* 107(8): 3864–3869.
- Raick X, Leccini D, Kever L, Colleye O, Bertucci F, Parmentier E (2018) Sound production mechanism in triggerfish (Balistidae): a synapomorphy. *Journal of Experimental Biology* 221(1): jeb168948.
- Ramírez-Rivera S, Bernal G (2019) Music is capable of inducing changes in gene expression in gastric cancer cells. *Journal of Gastrointestinal Cancer* 50(1): 175–180.
- Roth A, Tschardtke H (1976) Ultrastructure of the ampullary electroreceptors in lungfish and Brachiopterygii. *Cell and Tissue Research* 173(1): 95–108.

- Russo C, Patanè M, Pellitteri R, Stanzani S, Russo A (2021) Prenatal music exposure influences weight, ghrelin expression, and morphology of rat hypothalamic neuron cultures. *International Journal of Developmental Neuroscience* 81(2): 151–158.
- Russo C, Russo A, Gulino R, Pellitteri R, Stanzani S (2017) Effects of different musical frequencies on NPY and Ghrelin secretion in the rat hypothalamus. *Brain Research Bulletin* 132(Jun): 204–212.
- Samraus HH, Hecker PA (1985) Zum Einfluss von Geräuschen auf die Milchleistung von Kühen. (On the influence of noise on the milk yield of cows). *Berliner und Münchener Tierärztliche Wochenschrift* 98: 298–302.
- Sanyal T, Palanisamy P, Nag TC, Roy TS, Wadhwa S (2013) Effect of prenatal loud music and noise on total number of neurons and glia, neuronal nuclear area and volume of chick brainstem auditory nuclei, field L and hippocampus: a stereological investigation. *International Journal of Developmental Neuroscience* 31(4): 234–244.
- Saraogi M, Geraghty RM, Hameed B, Rob S, Pietropaolo A, Sarica K, et al. (2020) Role of complementary medicine (music, acupuncture, acupressure, TENS and audio-visual distraction) in shockwave lithotripsy (SWL): a systematic review from EAU sections of urolithiasis (EULIS) and uro-technology (ESUT). *Urology* 145(Nov): 38–51.
- Sarvaiya N, Kothari V (2015) Effect of audible sound in form of music on microbial growth and production of certain important metabolites. *Microbiology* 84(2): 227–235.
- Savage PE, Loui P, Tarr B, Schachner A, Glowacki L, Mithen S, et al. (2020) Music as a coevolved system for social bonding. *Behavioral and Brain Sciences* 44(Aug): e59.
- Sheikhi S, Saboory E (2015) Neuroplasticity changes of rat brain by musical stimuli during fetal period. *Cell Journal* 16(4): 448–455.
- Schöner MG, Simon R, Schöner CR (2016) Acoustic communication in plant-animal interactions. *Current Opinion in Plant Biology* 32(Aug): 88–95.
- Shah A, Raval A, Kothari V (2016) Sound stimulation can influence microbial growth and production of certain key metabolites. *Journal of Microbiology, Biotechnology and Food Sciences* 9(5): 330–334.
- Shaobin G, Wu Y, Li K, Ma S, Wang Q, Wang R (2010) A Pilot study of the effect of audible sound on the growth of *Escherichia coli*. *Colloids and Surfaces B: Biointerfaces* 78(2): 367–371.
- Suga N, O'Neill WE (1979) Neural axis representing target range in the auditory cortex of the mustache bat. *Science* 206(4416): 351–353.
- Syrovatkina V, Alegre KO, Dey R, Huang XY (2016) Regulation, signaling, and physiological functions of G-proteins. *Journal of Molecular Biology* 428(19): 3850–3868.
- Uchiyama M, Jin X, Zhang Q, Amano A, Bashuda H, Niimi N (2012a) Auditory stimulation of opera music induced prolongation of murine cardiac allograft survival and maintained generation of regulatory CD4<sup>+</sup>CD25<sup>+</sup> cells. *Journal of Cardiothoracic Surgery* 7(Mar): 26.
- Uchiyama M, Jin X, Zhang Q, Amano A, Watanabe T, Niimi, M (2012b) Music exposure induced prolongation of cardiac allograft survival and generated regulatory CD4<sup>+</sup> cells in mice. *Transplantation Proceedings* 44(4): 1076–1079.
- Usui C, Kirino E, Tanaka S, Inami R, Nishioka K, Hatta K, et al. (2020) Music intervention reduces persistent fibromyalgia pain and alters functional connectivity between the insula and default mode network. *Pain Medicine* 21(8): 1546–1552.
- Vannoni D, Albanese A, Battisti E, Aceto E, Giglioni S, Corallo C, et al. (2012) *In vitro* exposure of human osteoarthritic chondrocytes to ELF fields and new therapeutic application of musically modulated electromagnetic fields: biological evidence. *Journal of Biological Regulators and Homeostatic Agents* 26(1): 39–49.

- Veits M, Khait I, Obolski U, Zinger E, Boonman A, Goldstein A, et al. (2019) Flowers respond to pollinator sound within minutes by increasing nectar sugar concentration. *Ecology Letters* 22(9): 1483–1492.
- Ventura C, Gullà D, Graves M, Bergonzoni A, Tassinari R, von Stietencron J (2017) Cell melodies: when sound speaks to stem cells. *CellR4* 5(2): e2331.
- Vicient CM (2017) The effect of frequency-specific sound signals on the germination of maize seeds. *BMC Research Notes* 10(1): 323–327.
- Wachi M, Koyama M, Utsuyama M, Bittman BB, Kitagawa M, Hirokawa K (2007) Recreational music-making modulates natural killer cell activity, cytokines, and mood states in corporate employees. *Medical Science Monitor* 13(2): CR57–70.
- Wallin NL, Merker B, Brown S (2000) *The origins of music*. Cambridge, MS, US: MIT Press.
- Wang M, Yi G, Gao H, Wu B, Zhou Y (2020) Music-based interventions to improve fibromyalgia syndrome: a meta-analysis. *Explore* 16(6): 357–362.
- Warner RW (2009) The anatomy of the syrinx in passerine birds. *Journal of Zoology* 168(3): 381–393.
- Wei JK, Wang WC, Zhai RW, Zhang YH, Yang SC, Rizak J, et al. (2016) Neurons differentiated from transplanted stem cells respond functionally to acoustic stimuli in the Awake Monkey brain. *Cell Reports* 16(4): 1016–1025.
- Wiltschko FR, Wiltschko W (2012) Magnetoreception. *Advances in Experimental Medicine and Biology* 739: 126–141.
- Whitlow WLA (1993) *The sonar of dolphins*. New-York: Springer-Verlag.
- Xiong X, Han L, Liu S, Miao J, Luo M, Xue M, et al. (2018) Music intervention improves spatial learning and memory and alters serum proteomics profiling in rats. *Journal of Neuroscience Research* 96(10): 1727–1736.
- Yang H, Luo Y, Hu Q, Tian X, Wen H (2021) Affiliations benefits in Alzheimer’s disease of sensory and multisensory stimulation. *Journal of Alzheimer’s Disease* 82(2): 463–484.
- Zhang QL, Xu N, Huang ST, Lin ZW, Chen LW, Cao H, et al. (2020) Music therapy for early postoperative pain, anxiety, and sleep in patients after mitral valve replacement. *Thoracic and Cardiovascular Surgery* 68(6): 498–502.
- Zimmermann U, Fermin C (1996) Shape deformation of the organ of Corti associated with length changes of outer hair cell. *Acta Oto-Laryngologica* 116(3): 395–400.